

The Supernova-Gamma Ray Burst Connection

Lifan Wang¹ and J. Craig Wheeler¹

¹Astronomy Department, University of Texas, Austin, Texas 78712;

lifan@tao.as.utexas.edu, wheel@astro.as.utexas.edu

Received _____; accepted _____

ABSTRACT

Supernovae 1998bw and its corresponding relativistically expanding radio source are coincident with the γ -ray burst source GRB 980425. We show that of six recent SN Ib/c for which an outburst epoch can be estimated with some reliability, four have radio outbursts and all are correlated in time and space with BATSE γ -ray bursts. The joint probability of all six correlations is 1.5×10^{-5} . No such correlation exists for SN Ia and SN II. The γ -ray energy associated with the SN/GRB events is $\sim 10^{46} - 10^{48}$ ergs if emitted isotropically. Economy of hypotheses leads us to propose that all γ -ray bursts are associated with supernovae and that the γ -ray burst events have a quasi-isotropic component that cannot be observed at cosmological distances and a strongly collimated and Doppler-boosted component that can only be seen if looking nearly along the collimation axis. Such collimation requires a high rate of occurrence perhaps consistent with a supernova rate. The collimated flow may be generated by core collapse to produce rotating, magnetized neutron stars. All core collapse events may produce such jets, but only the ones that occur in supernovae with small or missing hydrogen envelopes, Type Ib or Ic, can propagate into the interstellar medium and yield a visible γ -ray burst. We suggest that asymmetries in line profiles and spectropolarimetry of SN II and SN Ib/c, pulsar runaway velocities, soft γ -ray repeaters and γ -ray bursts are associated phenomena.

Subject headings: supernovae: general – gamma rays: bursts – pulsars: general
– ISM: jets and outflows

1. Introduction

The first few months of 1997 brought a major breakthrough in one of the outstanding mysteries of modern astrophysics, the cosmic γ -ray bursts (Klebesadel, Strong & Olson 1973, Fishman 1995). The quest for an explanation of γ -ray bursts had been handicapped by a lack of direct knowledge of the distance to the bursts. This changed on February 28, 1997, when BeppoSAX localized a burst, GRB 970228, sufficiently well that an optical follow-up was feasible (Frontera et al. 1998). The result was the discovery of the first optical counterpart (van Paradijs et al. 1997). Two months later, in early May, BeppoSAX found another γ -ray burst enabling another optical identification (GRB 970508; Bond 1997). In this case, foreground absorption lines show that the source is at a redshift in excess of $z = 0.863$, proving that the source is at a cosmological distance (Metzger et al. 1997). These two optical counterparts have revolutionized the field and proven the power of focusing optical astronomy on this decades-old problem.

Early estimates of the energy of γ -ray bursts assuming cosmological distances gave energies comparable to the kinetic energy of supernovae, $\sim 10^{51}$ ergs, entirely in γ -rays. From the lower limit to the redshift of the optical counter-part of GRB 970508 and the observed maximum brightness, $R \approx 19.6$ mag (Castro-Tirado et al. 1997), one can deduce a distance and hence the absolute brightness. For isotropic emission, this event was about a factor of 100 brighter than a Type Ia supernova at a redshift of 1. A third optical counterpart was associated with GRB 971214. The apparent host galaxy for this event has a redshift of $z = 3.42$ (Kulkarni et al. 1998a). For isotropic emission, this event requires 3×10^{53} ergs.

2. SN 1998bw = GRB 980425

Soffitta et al. (1998) reported the detection by the BeppoSAX Gamma-Ray Burst Monitor of a γ -ray burst on Apr. 25.90915 UT (GRB 980425). The BATSE experiment on CGRO confirmed the detection (Kippen et al. 1998). The event was average in terms of the BATSE burst flux/fluence distribution. Galama et al. (1998) reported supernova 1998bw offset from the nucleus of the face-on barred spiral galaxy ESO 184-G82. This position is within the error box of GRB 980425, but does not coincide with either of the two X-ray sources of the BeppoSAX NFI image (Pian, et al. 1998). Lidman, et al. (1998) reported peculiar spectral properties for the supernovae. Wieringa et al. (1998) reported no radio sources were detected within the 1' error radius of the two NFI sources, but that there is a bright radio source that coincides with the optical astrometric position given by Galama et al. (1998). At the distance to ESO 184-G82, the radio source was more than three times as luminous as SN 1988Z, one of the most luminous radio supernovae. Given the seemingly small likelihood of finding the radio source and the supernova in this field, it is very probable that these events are associated with GRB 980425 despite the apparent discrepancy of the positions with the NFI locations. This is supported by analysis of the radio source that strongly suggests that it is expanding relativistically (Chevalier 1998, Kulkarni et al. 1998b). Independent of the association with the γ -ray burst, this is direct evidence that supernovae can produce relativistic outflows.

The association of a supernova with GRB 980425 may substantially alter the interpretation of the GRB afterglow. The current fireball model involves the shock interaction between relativistic ejecta and the CSM/ISM. The association with a supernova means that this interaction is *not* the only mechanism that produces optical emission although the interaction is certainly important for producing the radio emission and perhaps the γ -rays and X-rays (Mészáros & Rees 1997a). The UV/optical emission of SN 1998bw

is likely to consist of two parts, one from the stellar ejecta and the other from synchrotron emission produced in accord with the popular fireball model.

We have observed something similar to SN 1998bw before. The optical spectra of SN 1998bw resemble those of SN 1997ef. SN 1997ef and SN 1998bw are the only two supernovae known for which high quality spectroscopy was obtained and yet they can not be easily classified under the current classification scheme. At early times, SN 1997ef showed broad emission/absorption features with FWHM about $50,000 \text{ km s}^{-1}$, and all the features evolved rapidly to the red at a rate of about $1500 \text{ km s}^{-1}\text{day}^{-1}$ during the first week of discovery (Garnavich et al. 1997). At later times, the spectrum of SN1997ef resembled a Type Ic supernova which is characterized by little evidence for hydrogen, helium, or the strong Si II line of Type Ia. Sadler et al. (1998) suggest an association of SN 1998bw with a Type Ib, but it is clear that early spectra did not resemble canonical SN Ib or SN Ic. Optical spectropolarimetry of SN 1997ef did not reveal polarization to an upper limit of 0.8% (Wang, Howell, Wheeler 1998).

3. Other Supernova - Gamma-Ray Burst Associations

We have investigated the correlation of supernovae with BATSE γ -ray burst sources. Table 1 shows a compilation of all recent supernovae that had sufficiently well-established light curves to define maximum light and hence to yield an estimate of the time of explosion. Column 1 gives the identification of the supernova; column 2, the spectral type; column 3, the estimated dates of the explosion using the same notation as for γ -ray bursts, year, month, and day; column 4, the right ascension; column 5, the declination; column 6, whether there was a reported radio counterpart; column 7, the probability of at least one random association with a γ -ray burst, given as $1 - \prod_{i=1}^N (1 - P_i)$, where P_i is the BATSE $3\text{-}\sigma$ error box centered on the γ -ray burst divided by 4π steradians for each

γ -ray burst that occurred in the supernova temporal window of column 3 and N is the total number of these error boxes. Where more than one number is given in column 7, the probabilities were calculated using the 3-, 2-, and 1- σ BATSE error boxes, respectively. The underlined numbers indicate γ -ray burst counterparts were found within the error boxes of the corresponding γ -ray bursts, whereas non-underlined numbers indicate no associations. For the events where there is an associated γ -ray burst, the burst identification is given in column 1, and the RA and declination of the γ -ray burst in columns 4 and 5.

Table 1 shows that of the 6 well-observed SN Ib/c since CGRO/BATSE was launched, 4 had observed radio outbursts and *all* were correlated with a γ -ray burst. Four SN Ib/c were found to be within the 2- σ error boxes of the associated γ -ray bursts. The individual probabilities of random associations range from 0.06 to 0.3. The joint probability that all 6 associations are by chance is 1.5×10^{-5} . The real probability of accidental association is probably much smaller since the γ -ray bursts tend to occur at the best estimate of the time of the supernova explosion. The chance association in each case with the error box of the particular γ -ray burst we have identified is about 10^{-3} and the joint probability of the 6 events is vanishingly small.

Table 1 also gives data on 7 Type Ia events that are expected to undergo thermonuclear explosion and hence not core collapse. There is *no* correlation of any γ -ray burst with any of these events. Table 1 also contains 3 Type II events, none of which is correlated with γ -ray bursts and 4 events like SN 1993J (referred to here as SN IIt). These events are thought to have only thin hydrogen envelopes and hence to be nearly bare cores like SN Ib/c. Of the SN IIt in Table 1, there are no apparent correlations with γ -ray bursts except with SN 1993J, the nearest and brightest. A correlation with SN 1996cb is excluded only by dint of the very small BATSE error box for GRB 961212.

To assess the probability of random association, we have applied exactly the same test

for each supernova, but artificially placed all the supernovae on the exact opposite side of the sky. Only three supernovae, SN 1997ef (peculiar Ic), SN 199bq (Ia), and SN 1998T (Ib/c) were found to be within the $3\text{-}\sigma$ error boxes of 3 γ -ray bursts that occurred at epochs corresponding to the temporal windows for the supernova explosions. Considering that the probability for each association by random is about 20%, this number of correlations is entirely consistent with chance in a sample of 21 supernovae.

SN 1997ef was discovered about 10 days before maximum on JD 10778. As shown in Table 1, it was within the $2\text{-}\sigma$ BATSE error box of GRB 971115 on JD 10768 (Garnavich, 1998) and within the $3\text{-}\sigma$ error box of GRB 971120 on JD 10772. At most one association is real. Interestingly, SN Ib/c 1997ei was discovered at about the same time in roughly the same area of the sky. SN 1997ei is found to be within the $3\text{-}\sigma$ error box of GRB 971120.

4. The Case for Strong Collimation

The connection between γ -ray bursts and supernovae may call for a re-interpretation of the physical nature of γ -ray bursts. From the γ -ray fluence of GRB 980425 and the recession velocity (assuming $H_0 = 65 \text{ km s}^{-1}\text{Mpc}^{-1}$) the total γ -ray energy is 7×10^{47} ergs. The γ -ray energy of the other relatively nearby events associated with SN Ib/c in Table 1 is $\sim 10^{46} - 10^{48}$ erg assuming spherical geometry. This is completely consistent with an origin in supernovae. The question arises as to the nature of the very distant, apparently very energetic γ -ray bursts. These could also be related to supernovae if a portion of the γ -ray energy is strongly collimated. In particular, the supernova/ γ -ray burst connection suggests that we are dealing with energies that are a fraction of a supernova energy, perhaps ten percent, not 100 or more times a supernova energy. Strong collimation, in turn, demands a much higher incidence rate. If, for example, about 10^{50} ergs were collimated into 0.1 percent of 4π steradians in the form of a highly relativistic jet, the equivalent energy would be

10^{53} ergs if sphericity were assumed. The required rate of events would be correspondingly higher by a factor of 10^3 , so γ -ray bursts would have to occur about once in 100 years per bright galaxy, not once per 10^5 years, the canonical number for γ -ray burst and one associated with neutron star coalescence rates (Narayan, Piran & Shemi 1991). This rate could be lower by more than two orders of magnitude if the redshift for γ -ray bursts is as high as $z \sim 6$, giving a larger volume and lower rate per bright galaxy. A frequency of once per 100 years is a conventional rate for supernovae.

5. Discussion

Table 1 strongly suggests that at least one component of the γ -ray bursts arise in supernovae. The data shows that normal SN II and SN Ia are unlikely to produce γ -ray bursts. Table 1 does suggest that SN Ib/c are important γ -ray burst producers. Woosely (1993; see also Fryer & Woosely 1998) anticipated the utility of bare stellar cores as a site for γ -ray bursts. Woosely’s “failed Type Ib” scenario involved accretion from a torus onto a central black hole and associated neutrino production with mild collimation affecting the energetics and event rates. Paczyński’s (1997) “hypernova” scenario is generically related by invoking a “failed Type Ib” that makes a black hole, but invokes a large magnetic field to yield a “dirty fireball” with relatively large energy and again mild collimation. With strong collimation, the energy could correspond to that of a supernova with proportional effect on the event rates (Shapiro 1998). We suggest that it is successful SN Ib/c that make γ -ray bursts.

If γ -ray bursts, both at high and low redshift, are indeed produced by SN Ib/c, highly collimated, relativistic jets are required to solve the energy problem for the distant bursts. The hypothesis that all core collapse events produce relativistic beams is consistent with the current data. We would expect that such beams are stopped by the massive envelopes of

SN II, but that strong asymmetries are generated. This is consistent with the observations that all SN II are polarized and that almost no SN Ia are (Wang et al. 1996) and that SN II are not associated with γ -ray bursts. For progenitors that have lost most or all of their envelopes, the relativistic beams may survive to impact on the surroundings and create the phenomena associated with γ -ray bursts. Thus all SN Ib/c may produce γ -ray bursts and all γ -ray bursts may be associated with SN Ib/c.

This interpretation requires that there are both isotropic and non-isotropic components to the γ -ray bursts and associated phenomena. The isotropic component may be neither collimated nor Lorentz boosted and beamed. The anisotropic component must be subject to both. The γ -ray energy associated with the SN Ib/c, $\sim 10^{46} - 10^{48}$ ergs, is extremely faint compared with the other BeppSAX γ -ray bursts for which we know something about their distances. It is also a minor fraction of the energy released during a supernova event. If 10^{50} ergs of γ -ray energy were collimated into a cone of 10^{-3} of 4π steradians and pointed towards the Earth, we would see an apparent isotropic γ -ray burst energy of 10^{53} ergs, or a fluence of about 10 ergs/cm² at the distance of SN 1998bw. This implies that we are seeing in GRB 980425 the isotropic component of the γ -ray flux. In some other direction it would have presumably been a factor of $\sim 10^5$ brighter. Such an event would be visible at redshift greater than 3. Since only 1000 supernovae are known, it is very likely that a strongly collimated event would have escaped direct detection in a subset of some types of supernovae. Likewise, these events would not be seen at cosmological distances unless we are closely in the line-of-sight of the beam. The collimation required in this picture, confinement to an opening angle of a few degrees, is not unprecedented. Jets of comparable collimation occur in many astronomical environments.

The ratio of the luminosities of the collimated, boosted component and the isotropic component may be related to the *apparent* energy ratios, assuming both components to be

isotropic. Comparing GRB 971214 (Kulkarni et al. 1998a) to GRB 980425 gives a ratio of $3 \times 10^{53}/7 \times 10^{47}$ or 4×10^5 . If this is a representative ratio, then for the same detection limits the volume sampled by the distant collimated, boosted component is about 3×10^8 times that of the nearby component. If the distant component is collimated into only $\sim 10^{-3}$ of 4π steradians, there should be about 3×10^5 distant, strongly collimated γ -ray bursts for every “local” isotropic event. This means that in the current BATSE catalog there should be very few local events. The BATSE catalog should thus be dominated by the distant, collimated events, in complete accord with the isotropy of that sample. On the other hand, we apparently do see a few local events, the SN Ib/c of Table 1. To make our hypothesis plausible, the luminosity ratio of the collimated, beamed component to the isotropic component must be closer to $\sim 10^4$. There may be a distribution of collimation and Lorentz factors and boosted luminosities and corresponding Malmquist bias so that GRB 971214 is not typical.

From the brightness temperature Chevalier (1998) and Kulkarni et al. (1998b) have deduced that the radio source associated with SN 1998bw is expanding relativistically. It is surprising that a supernova produced relativistic ejecta. If the radio flux is reduced by a factor of $\sim 10^4$, one gets the characteristic radio flux for SN Ib/c. The radio emission from SN 1998bw could be from the jet, but with a viewing angle greater than zero, or perhaps after the jet has slowed and the radio emission is more nearly isotropic so there is less boost. A component of the radio could arise from more normal radio emission associated with collision of supernova ejecta with the circumstellar medium. We note that when it initially forms, the jet from the supernova core might be unstable, yielding beaming in time-dependent directions whereas the radio arises in a later blast wave phase when the propagation and beaming directions are well established. At the distance of SN 1998bw, radio interferometry should be able to resolve the jet and give some clues to its structure and anisotropy.

Spectropolarimetry of supernovae has already given evidence for strong anisotropy of the core collapse process. As remarked above, all SN II and SN Ib/c are polarized, nearly all SN Ia are not (Wang et al. 1996). There could be a number of reasons why the ejecta of supernovae are asymmetric, but the hypothesis that the core collapse process itself is a major contributor yields the prediction that for smaller buffering envelopes, the polarization should be greater. Indeed, SN Ic 1997X produced the largest polarization of any supernova yet measured, $\sim 6 - 7$ percent (Wang, Wheeler & Höflich 1998).

Core collapse supernovae may thus be highly anisotropic. This may be seen in the observations of SN 1997ef and SN 1998bw. From the width and temporal evolution of spectral features, SN 1997ef showed very high velocities at early times and a spectrum unlike other identified supernovae (Garnavich et al. 1997). This may mean that it was observed more nearly on axis. Polarization should be maximized when an event is viewed normal to the elongation axis and, indeed, SN 1997ef showed rather little polarization (Wang, Howell & Wheeler 1998). SN 1997ef presumably cannot have been beamed directly at us or we would have seen a very much brighter γ -ray source. The hydrodynamics and the γ -ray beam could plausibly have different opening angles. We would predict that SN 1998bw would also have relatively small polarization since the radio emission and the optical spectra suggest it was observed relatively pole-on.

If all core collapse events produce jets containing $\sim 10^{50}$ ergs of energy, about 10^{-3} of the neutron star binding energy, then current calculations of core collapse may need to be revised. Core collapse must probably include rotation and magnetic fields as intrinsic components. The question of how to make a tightly collimated jet from core collapse will clearly require separate effort. We note two relevant issues. Pulsars show typical runaway velocities of 500 km s^{-1} , requiring a kinetic energy of about 10^{49} erg. Pulsar runaway might well be linked to the momentum imbalance involved in generating two not quite equal jets

with total energy of about 10^{50} ergs. The formation of jets might also be enhanced if the magnetic field were larger than often assumed. Models for soft γ -ray repeaters require fields of $\sim 10^{15}$ Gauss (Duncan & Thompson 1992). Recent detection of a spin down in a soft γ -ray repeater is consistent with such a large field (Kouveliotou et al. 1998). The calculation of LeBlanc and Wilson (1970) of a rotating, magnetic core yielded a magnetic field of $\sim 10^{15}$ gauss and an axial jet of with an energy of $\sim 10^{50}$ ergs. Blackman & Yi (1998; see also Thompson, 1994; Mészáros & Rees 1997b) have noted that the energy in a jet associated with a pulsar might be in the form of large amplitude electromagnetic waves, rather than relativistic particles, *per se*.

By the hypothesis presented here, SN 1987A must have created an embryonic jet in its core. Even if this hypothesis is correct, it is not clear how such a jet could penetrate the hydrogen envelope, which is known to have been massive. Nevertheless, it might be appropriate to reexamine the “mystery spot” in SN 1987A.

The authors are grateful to Peter Diener, Rob Duncan, and Ed Robinson for helpful discussions and to the Crown & Anchor for hospitality as the ideas presented here were developed. This research was supported in part by NSF Grant 95-28110, a grant from the Texas Advanced Research Program, and by NASA through grant HF-01085.01-96A from the Space Telescope Science Institute which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS 5-26555.

REFERENCES

- Blackman, E. G. & Yi, I. 1998, *ApJ*, 498, L31
- Bond, H. E. 1997, *IAUC*, 6654
- Castro-Tirado, A. J. et al. 1997, *IAUC*, 6657
- Chevalier, R. A. 1998, private communication
- Duncan, R. C. & Thompson, C. 1992, *ApJ*, 392, L9
- Fishman, G. J. 1995, *PASP*, 107, 1145
- Frontiera, F. et al. 1998, *ApJ*, 493, L67
- Fryer, C. L. & Woosely, S. E. 1998, *astro-ph/9804167*
- Galama, T. J. et al. 1998, *IAUC* 6895
- Garnavich, P., Jha, S., Kirshner, R., Challis, P., Balam, D. Brown, W. & Bricena, C. 1997, *IAUC* 6786
- Garnavich, P. 1998, private communication
- Kippen, R. M. et al. 1998, *GCN* 67
- Klebesadel, R. W., Strong, I. B. & Olson, R. A. 1973, *ApJ*, 182, L85
- Kouveliotou, C. et al. 1998, *ApJ*, in press
- Kulkarni, S. et al. 1998a, *Nature*, 393, 35
- Kulkarni, S. et al. 1998b, *IAUC* 6903
- LeBlanc, J. M. & Wilson, J. R. 1970, *ApJ*, 161, 541
- Lidman, C. et al. 1998, *IAUC* 6895
- Mészáros, P. & Rees, M. J. 1997a, *ApJ*, 476, 232
- Mészáros, P. & Rees, M. J. 1997b, *ApJ*, 482, L29

- Metzger, M. R. et al. 1997, *Nature*, 387, 878
- Narayan, R., Piran, T. & Shemi, A. 1991, *ApJ*, 379, L17
- Paczynski, B. 1998, *ApJ*, 494, L45
- Pian, E. et al. 1998, *GCN* 61
- Sadler, E. M., Stathakis, R. A., Boyle, B. J. & Ekers, R. 1998, *IAUC* 6901
- Shapiro, P. 1998, private communication
- Soffitta, P. et al. 1998, *IAUC* 6884
- Thompson, C. 1994, *MNRAS*, 270, 480
- van Paradijs, J. et al. 1997, *Nature*, 386, 686
- Wang, L., Howell, D. A., & Wheeler, J. C. 1998, *IAUC*, 6820
- Wang, L., Wheeler, J. C., Li, Z. W., & Clocchiatti, A. 1996, *ApJ*, 467, 435
- Wang, L., Wheeler, J. C. & Höflich, P. 1998, in *SN 1987A: Ten Years After* ed. M. M. Phillips and N. Suntzeff (Provo: Ast. Soc. of the Pacific), in press
- Wieringa, et al. 1998, *IAUC* 6896
- Woosley, S. E. 1993, *ApJ*, 405, 273

Table 1: Supernovae and GRB Association

SN/GRB	Type	Dates	RA (2000.0)	DEC (2000.0)	Radio	Prob.
SN 1992H	II	920204-930211	13 56 19.6	+47 14 34.1	No	14.8
SN 1992ad	Ib	920609-920605	12 26 49.6	+08 52 38.7	Yes	
GRB 920609			13 06 19.2	-05 34 48.0		<u>29.8,14.0,1.7</u>
SN 1993J	II _t	930323-930328	09 55 24.8	+69 01 13.7	Yes	
GRB 930324			12 00 48.0	+59 34 12.0		<u>7.7,3.5,0.9</u>
SN 1994D	Ia	940328-940307	12 34 02.3	+07 42 05.7	No	16.2
SN 1994I	Ib/c	940329-940402	13 29 54.1	+47 11 30.5	Yes	
GRB 940331			10 31 14.4	+57 31 48.0		<u>8.1,3.7</u>
SN 1995D	Ia	950131-950109	09 40 54.8	+05 08 26.2		22.3
SN 1996B	II _t	951219-960109	12 23 56.6	+48 46 42.0		10.9
SN 1996N	Ib	960210-960224	03 38 55.4	-26 20 04.0	Yes	
GRB 960221			03 11 04.8	-31 14 24.0		<u>5.8,2.6,0.7</u>
SN 1996W	II	960402-960411	11 59 30.1	-19 16 05.0	No	6.2
SN 1996X	Ia	960315-960329	13 18 01.1	-26 50 45.3	No	18.9
SN 1996cb	II _t	961208-961215	11 03 42.0	+28 54 13.7	Yes	2.4
SN 1997ef	?	971113-971125	07 57 02.8	+49 33 40.2		
GRB 971115			05 38 31.2	+41 41 24.0		<u>28.0,13.4,3.5</u>
GRB 971120			10 23 24.0	+76 24 36.0		<u>28.0,13.4</u>
SN 1997X	Ib/c	961231-970121	12 48 14.3	-03 19 58.5	Yes	
GRB 970103			11 56 19.2	-06 51 00.0		<u>13.8,6.35</u>
SN 1997ei	Ic	971103-971203	11 54 59.0	+58 29 26.4		
GRB 971120			10 23 24.0	+76 24 36.0		<u>36.1,17.8,4.7</u>
SN 1997bp	Ia	970316-970401	12 46 53.8	-11 38 33.2		16.3
SN 1997bq	Ia	970401-970408	10 17 05.3	+73 23 02.1		21.1
SN 1997br	Ia	970320-970409	13 20 42.4	-22 02 12.3		46.8
SN 1997dd	II _t	970730-970813	16 05 46.0	+21 29 14.0		16.4
SN 1998S	II _n	980216-980302	11 46 06.0	+47 29 00.0		17.8
SN 1998T	Ib	980208-980303	11 28 33.2	+58 33 43.7		
GRB 980218			12 13 09.6	+56 38 24.0		<u>21.2,10.0</u>
GRB 980223			14 45 09.6	+28 12 36.0		<u>21.2,10.0</u>
SN 1998bu	Ia	980428-980503	10 46 46.0	+11 50 07.5		0.197